


Accepted Manuscript

Classification schemes for carcinogenicity based on hazard-identification have become outmoded and serve neither science nor society

Alan Boobis, Samuel Cohen, Vicki Dellarco, John E. Doe, Penelope A. Fenner-Crisp, Angelo Moretto, Timothy P. Pastoor, Rita Schoeny, Jennifer Seed, Douglas C. Wolf



PII: S0273-2300(16)30303-8

DOI: 10.1016/j.yrtph.2016.10.014

Reference: YRTPH 3706

To appear in: *Regulatory Toxicology and Pharmacology*

Received Date: 13 October 2016

Revised Date: 20 October 2016

Accepted Date: 21 October 2016

Please cite this article as: Boobis, A., Cohen, S., Dellarco, V., Doe, J.E., Fenner-Crisp, P.A., Moretto, A., Pastoor, T.P., Schoeny, R., Seed, J., Wolf, D.C., Classification schemes for carcinogenicity based on hazard-identification have become outmoded and serve neither science nor society, *Regulatory Toxicology and Pharmacology* (2016), doi: 10.1016/j.yrtph.2016.10.014.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Classification Schemes for Carcinogenicity Based on Hazard-identification Have Become Outmoded
and Serve neither Science nor Society**

**Alan Boobis^a, Samuel Cohen^b, Vicki Dellarco^c, John E Doe^d, Penelope A Fenner-Crisp^e, Angelo
Moretto^f, Timothy P Pastoor^g, Rita Schoeny^h, Jennifer Seedⁱ and Douglas C Wolf^j.**

a. Centre for Pharmacology & Therapeutics, Toxicology Unit, Department of Medicine,
Hammersmith Campus, Imperial College London, London, W12 0NN, UK

b. Department of Pathology and Microbiology, Havlik-Wall Professor of Oncology, University of
Nebraska Medical Center, Omaha, NE 68198-3135, USA

c. Independent consultant, Silver Spring, MD 20901, USA

d. Parker Doe LLP, Carpenter Court, Maple Road, Bramhall, Stockport, Cheshire, SK7 2DH, UK

e. Independent Consultant, North Garden, VA 22959, USA

f. Dipartimento di Scienze Biochimiche e Cliniche (Department of Biomedical and Clinical Sciences)
Università degli Studi di Milano, Milan, Italy

g. Pastoor Science Communication, LLC, Greensboro, NC, 27455, USA

h. Rita Schoeny LLC, Washington DC 20002

i. Independent Consultant, Alexandria, VA 22301, USA

j. Syngenta Crop Protection, LLC; Greensboro, NC, USA, 27419

Corresponding Author: John Doe – john.doe@parkerdoe.com

Abstract

Classification schemes for carcinogenicity based solely on hazard-identification such as the IARC monograph process and the UN system adopted in the EU have become outmoded. They are based on a concept developed in the 1970s that chemicals could be divided into two classes: carcinogens and non-carcinogens. Categorization in this way places into the same category chemicals and agents with widely differing potencies and modes of action. This is how eating processed meat can fall into the same category as sulfur mustard gas. Approaches based on hazard and risk characterization present an integrated and balanced picture of hazard, dose response and exposure and allow informed risk management decisions to be taken. Because a risk-based decision framework fully considers hazard in the context of dose, potency, and exposure the unintended downsides of a hazard only approach are avoided, e.g., health scares, unnecessary economic costs, loss of beneficial products, adoption of strategies with greater health costs, and the diversion of public funds into unnecessary research. An initiative to agree upon a standardized, internationally acceptable methodology for carcinogen assessment is needed now. The approach should incorporate principles and concepts of existing international consensus-based frameworks including the WHO IPCS mode of action framework.

Highlights

- Cancer classification schemes based on hazard-identification such IARC and UN GHS are outmoded and inadequate to manage chemical risks
- Chemicals with many orders of magnitude difference in potency and modes of action are placed in the same category
- Unintended consequences of unnecessary health scares, economic costs, loss of beneficial products, and diversion of public funds

- Globally accepted problem formulation and hypothesis-based frameworks provide modern approaches based on hazard and risk characterization
- Call for an international initiative to develop a consensus on classification methodology for carcinogenicity assessment

Key Words: Classification; Hazard Characterization; Risk Assessment; Carcinogenicity; IARC; GHS

Abstract word count: 199

Text Word Count: 7182

References Word Count: 1610

Contents

1. Introduction

2. Advances in Public Health and Chemical Risk Management

3. Classification and Risk Assessment

4. Problems with Classification

5. Cancer as a Major Concern

6. Cancer Classification

7. Introduction of the Cancer Bioassay

8. Increasing Understanding of Carcinogenicity (and its Consequences on Risk Assessment and Management)

9. Unchanged Processes Become Outmoded

10. Problems Resulting From Use of Outmoded Processes

11. Problem Formulation: What Problem is Being Addressed?

12. Reproductive Toxicology and Endocrine Disruption

13. Conclusions

69 Abbreviations

70 ACGIH – American Conference of Government Industrial Hygienists

71 CoC – United Kingdom Committee on Carcinogenicity

72 ECHA - European Chemicals Agency

73 EFSA – European Food Safety Authority

74 EPA – United States Environmental Protection Agency

75 EU – European Union

76 GHS - United Nations Global Harmonized System for Classification and Labelling

77 IARC – International Agency for Research on Cancer

78 IPCS – International Programme on Chemical Safety

79 JMPR –Joint FAO/WHO Meeting on Pesticide Residues

80 MOA – Mode of Action

81 NCI – United States National Cancer Institute

82 PMRA - Health Canada Pest Management Regulatory Agency

83 WHO – World Health Organization

84

1. Introduction

Cancer prevention is the primary objective of the evaluation of chemicals for their human carcinogenicity potential. This objective, however, is undermined by confusion resulting from conflicting pronouncements coming from multiple international and national agencies (Guardian, 2016). This has led to carcinogen definition and regulation being called “the poor relation to other cancer preventative measures” (Lancet, 2016). The problem arises from the different concepts and approaches that are being used, some of which were developed half a century ago. Their appropriateness was questionable at the time and they have now clearly become out of step with advances in scientific understanding and modern regulatory science.

Classifying chemicals on hazard-identification alone is one such outmoded concept. The International Agency for Research on Cancer (IARC) classification process for carcinogenicity and the United Nations Global Harmonized System for Classification and Labelling (GHS) (adapted and adopted in the EU and elsewhere) processes for carcinogenicity (and reproductive toxicity) are based on this outmoded concept.

The original intention of these processes was to raise a warning flag for chemicals of potential concern which would lead to fuller evaluation to determine if risk management measures need to be taken. However, the warning flags are never removed, and sometimes they even appear after more complete evaluation by regulatory authorities has determined that adequate risk management is in place. Of even greater concern is that evaluation often stops at classification and acceptability is based only on hazard with no consideration of the potential risk under even extreme (though remotely possible) human exposure.

This hazard-identification only process places chemicals with widely differing potencies and very different modes of action into the same category. Processed meat (consumption) and sulfur mustard gas are placed into the same category (group 1) by IARC as described in section 6. This leads

to confusion; should we treat processed meat as we do sulfur mustard gas – reduce exposure to zero; or should we treat sulfur mustard gas as we do red meat – consider it part of a healthy life style in moderation? This categorization can thus lead to unnecessary public anxiety; resources may be diverted that would be better used addressing more substantial problems; safe and useful products come under unnecessary and excessive scrutiny; and they may even be replaced by other less characterized and potentially less safe products.

This present work describes the origins of classification schemes based on hazard-identification, acknowledges that they were once useful, explains why they no longer serve a useful role and illustrates how science-based approaches in a risk based decision framework are more suited to protecting human health in the 21st century.

2. Advances in Public Health and Chemical Risk Management

The 20th century saw great advances in the state of public health; managing the potential risks from chemicals has played its part. Life expectancy increased by over 30 years in Europe and the Americas between 1900 and 2000 (Roser, 2015). Certain chemicals and technologies developed in the late 19th century and early 20th century did come at a price, however. At the time, there was poor understanding of the range of biological effects that chemicals could cause until the pioneering observational studies that identified how chemicals could adversely affect human health were published (Goldblatt, 1944). Many adverse effects observed in humans were then verified in animal studies. By the middle of the 20th century there was a shift towards the use of animal studies to predict what could happen in humans, which led, in the 60's and 70's, to the development of extensive and diverse toxicological studies to identify and characterize chemical hazards, and predict the human safe dose, before adverse effects could occur in humans. Hazard-identification and characterization via animal studies became the standard for predicting and then avoiding potential

adverse effects in humans. As a result of this approach to chemical safety assessment, exposure to high-risk chemicals has been progressively reduced (Kauppinen et al, 2013). Whilst not perfect, this approach has the advantage that chemicals potentially toxic to humans are identified before there is any human exposure.

3. Classification and Risk Assessment

The results of laboratory animal toxicology studies are used for identifying in animals adverse health effects assumed without additional information to represent a potential hazard to humans which may be further characterized in terms of severity and dose response. This information is then most appropriately used for assessing potential human health impact from the use or presence in the environment of the chemical. There are two major ways in which this is done: risk assessment and classification.

Risk assessment requires estimation of the human exposure in terms of duration, frequency and magnitude to derive a plausible maximum dose to which humans might be exposed. This dose is then compared with the projected safe human dose level derived from hazard characterization; if the projected exposure is lower than the projected human safe dose then safety in use can be assumed, and if not then it may be necessary to identify and implement risk mitigation measures. Risk assessment also requires evaluation of the relevance of the findings at high doses in animal studies to lower exposures in humans. Mechanisms leading to toxicity in animals might not be relevant to humans, or changes occurring at high doses might not be relevant to low doses. In other words, scientific evaluations are necessary.

Classification uses a different approach while being based on similar principles. It focuses on the hazard which has been identified, usually from animal studies and, then, grades the hazard into

various categories based on the severity and, in some instances, dose response. Classification was originally intended to provide information on the effects of a chemical following acute exposure for labelling purposes for transport (UN, 2011). However, its use has broadened substantially so that many regulatory schemes are based solely on classification for a range of end points following either acute or repeated exposure leading directly to risk management action without consideration of the chemical potency, severity of the effect or mode of action or the nature and extent of human exposure.

4. Problems with Classification

The advantages and disadvantages of both approaches have been reviewed by Barlow et al, 2015, who concluded that both approaches have their uses depending on the situation being addressed. Classification is more appropriate for acute toxicity or in situations where it is hypothesized that there is no threshold for an adverse effect. It requires less data and can be valuable in providing guidance when a decision has to be taken before a full evaluation has been carried out. Risk assessment provides more information and insight into the magnitude of risks, and can be used as a basis for deriving “safe” levels of exposure. However, problems can arise when both approaches are used in regulation by the same or different agencies that address the same agent/substance. This separation of decision-making can result in hazard-based restrictions on marketing and use or unnecessary remediation of environmental levels, even when risk-based assessments show there is reasonable certainty no harm will result. This in turn can lead to contradictory, confusing and ultimately unnecessary actions.

These problems arise most often when the classification process focuses simply on identifying the hazard but does not go on to characterize the hazard in terms of severity, dose response and mode

of action. This is the situation with some schemes in the areas of carcinogenicity, and reproductive toxicity, and it is a source of the current controversy on how to prioritise and manage the risk posed by chemicals identified as endocrine disrupters. The omission of characterization of both hazard and risk in these schemes stems from the historical special treatment given to certain health effects based on their public perception (Slovic, 1987) such as carcinogenic and reproductive toxicity effects, and more recently endocrine disruption, neurotoxicity and immunotoxicity.

5. Cancer as a Major Concern

Cancer had become such a concern that in 1971 US President Nixon signed the National Cancer Act saying in his State of the Union Address “The time has come when the same type of concentrated effort that split the atom and took a man to the moon should be turned toward conquering this dread disease”. Large sums of money were devoted to reducing cancer deaths and prevention was seen as part of the “war on cancer” as well as the discovery and development of treatments. It was believed, at the time, that a large proportion of cancers were caused by industrial chemicals. Hueper (1955) from the National Cancer Institute (NCI) had concluded that cancer from exposure to industrial chemicals was of far greater concern than cancer from tobacco smoking. This has since been shown to be incorrect as tobacco smoking is now recognized as the second leading cause of death globally, with 6.3 million deaths annually attributed to smoking (GBD2010).

In the middle of the 20th century, the concept developed that chemicals could be segregated into two classes: carcinogens and non-carcinogens. It was postulated that a major reduction in cancer incidence would result if we could identify the “carcinogens” and, hence, replace them with “non-carcinogens”. This concept started the drive towards using hazard-identification alone for carcinogenicity which has continued for nearly half a century in a largely unmodified way. It was also the basis for the use of the Maximum Tolerated Dose in animal studies which was thought to optimize the chances of identifying “carcinogens”. Whilst the concept was considered sound in

principle at the time, it was based on a fundamental misconception. There is now a greater understanding of the complex biology and etiology of cancer, specifically how chemical exposure can lead to cancer, and the idea of a binary separation into “carcinogens” and “non-carcinogens” has proved to be overly simplistic. Indeed, a very wide range of chemical cans cause cancer under the “right” experimental circumstances many of which having no relevance to humans or achievable exposure levels (see section 7).

6. Cancer Classification

Carcinogen hazard-identification is primarily based on the evaluation of human epidemiological data, if available, and the results of long term bioassays in laboratory rodents. At first most of the evaluations were based upon human epidemiology studies in occupational settings. Processes were set up by several national and international bodies to identify carcinogens that were largely based on Sir Austin Bradford Hill’s considerations for causality (Hill, 1965). The strength of the evidence for causality varied and, therefore, it was graded to allow chemicals to be placed into different categories regarding the confidence in a causal link for carcinogenicity in humans.

The categories used today by IARC (2012) still reflect this:

- Group 1: The agent is *carcinogenic to humans*
- Group 2A: The agent is *probably carcinogenic to humans*
- Group 2B: The agent is *possibly carcinogenic to humans*.
- Group 3: The agent is *not classifiable as to its carcinogenicity to humans*
- Group 4: The agent is *probably not carcinogenic to humans*.

Whether by accident or design, this was a classification system and, therefore, it was not surprising that this system was co-opted into some of the chemical classification schemes which were emerging in the 1970s and 1980s. However, unlike the other classification systems developed for

health protection, the carcinogenicity scheme deliberately avoided the valuable context of hazard or of dose response, severity, and mode of action or exposure. At the time this scheme was put in place, such characterization was not thought necessary as the aim was simply to identify “carcinogens” and to eliminate them.

The EU in its Classification and Labelling Guidelines (ECHA 2012) has implemented the UN GHS categorization system which is very similar using strength of evidence involving only the enumeration of tumors in human and animal studies and determination of their level of statistical significance. The Guidelines state that *“Sufficient human evidence demonstrates causality between human exposure and the development of cancer, whereas sufficient evidence in animals shows a causal relationship between the substance and an increased incidence of tumours. Limited evidence in humans is demonstrated by a positive association between exposure and cancer, but a causal relationship cannot be stated. Limited evidence in animals is provided when data suggest a carcinogenic effect, but are less than sufficient. The terms ‘sufficient’ and ‘limited’ have been used here as they have been defined by the International Agency for Research on Cancer (IARC).”*

The UN GHS categories are

- Category 1A: Known to have carcinogenic potential for humans, the placing of a substance in this category is largely based on human evidence
- Category 1B: Presumed to have carcinogenic potential for humans: the placing of a substance in this category is largely based on animal evidence
- Category 2: Suspected Human carcinogen

Thus a system set up half a century ago based on an overly simplistic concept as an initial attempt to address the disease burden of cancer has found its way into classification schemes for chemicals and also into some downstream risk management processes. The consequences may not have been

intended but they were predictable. The pressure to replace a chemical which had been identified as a potential human carcinogen was immense, even on the basis of animal studies, and the very act of categorizing a chemical in this way leads to a stigma which would often result in major changes in its use, including withdrawal, whereas risk-based assessments show that there is reasonable certainty no harm will result from its use.

7. Introduction of the Cancer Bioassay

All of this was occurring at the time as the potential of chemicals to cause adverse effects was being recognized and animal models were being developed. The long term bioassay for carcinogenicity was accepted as an OECD guideline study in 1981 (OECD, 2009) and the results of rodent bioassays were used as evidence, either alongside or instead of epidemiology, in deciding whether a chemical should be classified as a “carcinogen”. Since known human carcinogens, based on epidemiology studies, also caused cancer in animal models, it was concluded that a chemical that caused cancer in an animal model must also cause cancer in humans. Their reverse incorrect logic has been proven wrong numerous times. For example, about 60% of pharmaceuticals tested in the rodent bioassay gave positive results, but have been deemed safe for human use (Brambilla et al., 2012). Applying a hazard classification would have kept these life-saving pharmaceuticals off the market, including statins and proton pump inhibitors, two of the most widely used classes of drugs today.

The bioassay was intended for hazard-identification and was therefore designed to maximize the ability to detect “carcinogenicity”. Dosing was for as much of the life time of the animals as possible. Historically, exposure started after weaning, some newer study designs start exposure before birth. The highest dose was set as a Minimally Toxic Dose that would not impact the animals’ normal lifespan from effects other than cancer. This evolved into the Maximum Tolerated Dose (MTD),

which increased the doses used in an attempt to increase sensitivity to detect the “carcinogenicity” of low potency compounds. Under these assay conditions 50% of chemicals, both synthetic and natural, were capable of increasing the incidence of neoplasms (both malignant and benign) (Gold et al 1989). Numerous studies have shown that as the experimental dose of a chemical is increased, different saturable or inducible toxicokinetic (e.g., metabolism, uptake, excretion) and toxicodynamic (e.g., homeostasis, receptor interactions, protein binding, repair mechanisms) processes involved in chemical toxicity (e.g., tumorigenicity) can be involved, which may not be engaged at environmental exposures (Slikker et al 2004a,b). It seems that the high doses used were triggering different mechanisms which lead to the development of neoplasms in laboratory animals. Determining which mechanisms are operative along the dose-response curve has important implications for interpreting bioassay data for the purposes of predicting human risk.

The classification processes were, therefore, adapted to include the results of the animal bioassays in determining the strength of evidence for carcinogenicity. Induction of excess neoplasms in rodents, irrespective of the dose, was taken as strong evidence for carcinogenicity in humans and chemicals were categorized accordingly. The large proportion of chemicals classified in this way led to questioning of the validity of the assays and the overall process (Ames and Gold 1990). The finding that around 50% of chemicals caused neoplasms in these assays undermines the concept of separating chemicals into “carcinogens” and “non-carcinogens”. It is not logical that half the chemicals in use cause cancer in humans particularly given the high experimental doses used and the high incidence of background tumors in certain rodent species and strains and the lack of confirmation in scores of human epidemiology studies (Pastoor and Stevens, 2005).

8. Increasing Understanding of Carcinogenicity (and its Impact on Risk Assessment and Management)

Increasing understanding of chemically-driven carcinogenic pathways over the last several decades has progressively raised questions regarding the relevance for human health of certain tumor findings in rodent bioassays. Given the issues and debates around the human relevance and dose response of rodent tumors, advances in knowledge of chemical carcinogenesis, and emerging cost and time-effective methods to investigate modes of action (MOAs), the international need for harmonised guidance on how to look at mode of action information in cancer assessment was recognized. Work under the auspices of the WHO International Programme for Chemical Safety (IPCS) began in the 1990's to develop a weight of evidence framework (Sonich-Mullen et al., 2001). In the first stage of the framework one determines whether it is possible to establish an MOA for the animal tumor(s) under investigation by identifying a series of key events along the causal pathway to cancer using a weight-of-evidence approach based on the Bradford Hill considerations. The key events are compared first qualitatively and then quantitatively between those which would occur in the experimental animals and those which would occur in humans. Finally, a clear statement of confidence, analysis, and implications for risk assessment is produced.

The resulting IPCS mode of action framework was an important development in moving cancer assessment away from a phenomenological approach and toward enabling the integration of a fuller biological understanding of how chemicals induce neoplasia and a better understanding of the dose response relationships. Shortly afterwards, the IPCS framework was expanded to address how MOA knowledge can be used to evaluate the human relevance of animal responses based on species concordance analyzes (Meek et al., 2003, Boobis et al., 2006). In the early mid 2000's, the approach was extended to evaluate non-cancer endpoints and life stage information (Boobis et al., 2008; Seed et al., 2005) and to incorporate the quantitative consideration of dose response (Julien et al, 2009; Simon et al, 2014). IPCS has now updated the framework to consolidate the international work that

had been done and to emphasize that MoA analyzes should be problem formulation-based recognizing that MOA knowledge can inform different risk management decisions: priority setting, read across, or guiding research, not just for risk assessment where it has been most frequently used (Meek et al.2014).

When using this approach for assessing carcinogenicity, there are three broad outcomes which have an impact on how the chemical should be further evaluated:

- Rodent carcinogens that are considered relevant for humans and which have mode(s) of action indicating that there is no presumption of a threshold for the dose response.
- Rodent carcinogens that are considered relevant for humans and which have a mode(s) of action indicating that there is a threshold for the dose response. These often result from modes of action associated with the high experimental treatment doses that result in secondary processes (e.g., sustained cytotoxicity and compensatory hyperplasia) where a no effect level or margin of safety can be established. A substantial number of chemicals have been shown to fall into this group including the pesticides acifluorfen sodium, amitrole, captan, cyproconazole, folpet, lactofen, and pyroxasulfone (EPA, 2015)
- Rodent carcinogens that are considered to have mode(s) of action not relevant for humans. A number of chemicals induce tumors by modes of action well documented to be non-relevant to humans. Some examples are: kidney tumors in male rats associated with substances causing α 2u-globulin nephropathy ;pheochromocytomas in male rats exposed to particulates through inhalation secondary to hypoxemia; Leydig cell adenomas induced by dopamine antagonists or gonadotropin-releasing hormone (GnRH); certain thyroid tumors in rodents mediated by UDP glucuronyltransferase (UGT) induction (listed in the Classification Guidelines by ECHA, 2013).

The process for the assessment of the carcinogenicity of chemicals by many regulatory authorities or organisations has incorporated the concept that different modes of action have different implications for human safety. Examples are:

US EPA - The US EPA (2005) revised their Cancer Risk Assessment Guidelines to bring in more relevant science in the cancer risk assessment process by incorporating a framework for analyzing mode of action (consistent with the WHO IPCS approach). The US EPA also replaced their cancer categories with descriptors and weight of evidence narratives, and to acknowledge that carcinogens should be considered in ways appropriate to their full hazard and risk characterization. In the absence of data, the US EPA takes a public health protective position that animal tumor findings are assumed to be relevant to humans, and cancer risks are assumed to conform with the default hypothesis of non-threshold, low dose linearity. However, sufficient, scientifically justifiable mode of action information can support different conclusions. Non-linear dose response modelling may be appropriate. In some cases, the animal tumors are concluded to be not relevant to humans and thus not to be used in human risk assessment. These considerations are reflected in the descriptors which the EPA applies. More than one descriptor can be used when an agent's effects differ by dose or exposure route. For example, an agent may be "Carcinogenic to Humans" by one exposure route but "Not Likely to Be Carcinogenic" by a route by which it is not absorbed. Also, an agent could be "Likely to Be Carcinogenic" above a specified dose but "Not Likely to Be Carcinogenic" below that dose because a key event in tumor formation does not occur below that dose. These are descriptors which enable the US EPA to apply the appropriate risk assessment methodology.

ACGIH - The American Conference of Government Industrial Hygienists (ACGIH, 2016) has developed categories which describe different classifications based on the concept that different modes of action have different implications for human safety:

- A1- Confirmed human carcinogen

- A2 - Suspected human carcinogen
- A3 - Animal carcinogen. The agent is not likely to cause cancer in humans except under uncommon or unlikely routes or levels of exposure. The agent is carcinogenic in experimental animals at a relatively high dose, by route(s) of administration, at site(s), of histologic type(s), or by mechanism(s) that may not be relevant to worker exposure.
- A4 - Not classifiable as a human carcinogen
- A5 - Not suspected as a human carcinogen

UK CoC - The United Kingdom Committee on Carcinogenicity (CoC, 2012) developed a decision tree approach which takes into account the mode of action in the way the hazard is characterized and the risk assessed before risk management decisions are taken. The decision tree reviews the carcinogenicity data and leads to one of three conclusions:

- Exposure should be as low as reasonably possible for substances with a genotoxic mode of action
- Exposure should be below a level set using identification of critical end points and use of uncertainty factors for substances with other modes of action considered relevant to humans.
- Exposure should be below levels determined by consideration of non-carcinogenicity end-points for substances with modes of action for carcinogenicity considered non-relevant to humans.

SCOEL - The EU Scientific Committee on Occupational Exposure Limits (SCOEL) set criteria to include mode of action and strength of available data to provide input to the management of carcinogens (Bolt and Huici-Montagud, 2008):

- A) Non-threshold genotoxic carcinogens; for low-dose assessment of risk, the linear non-threshold (LNT) model appears appropriate. For these chemicals, regulations (risk

management) may be based on the ALARA principle ("as low as reasonably achievable"), technical feasibility, and other socio-political considerations.

B) Genotoxic carcinogens, for which the existence of a threshold cannot be sufficiently supported at present. In these cases, the LNT model may be used as a default assumption, based on the scientific uncertainty.

C) Genotoxic carcinogens with a practical threshold, as supported by studies on mechanisms and/or toxicokinetics; health-based exposure limits may be based on an established NOAEL (no observed adverse effect level).

D) Non-genotoxic carcinogens and non-DNA-reactive carcinogens; for these compounds a true ("perfect") threshold is associated with a clearly founded NOAEL.

Each of these modern schemes derives the dose which is predicted to be of concern, or not, in humans based on the concept that different modes of action have different implications for human safety.

9. Unchanged Processes Become Outmoded

Although it has not modified its categories since 1971, IARC currently makes no claim that its role is anything other than hazard-identification; *"These categories refer only to the strength of the evidence that an exposure is carcinogenic and not to the extent of its carcinogenic activity (potency)."* (IARC, 2015a). However, on occasion, IARC has pronounced on the risk of some of the carcinogenic hazards that they have identified (IARC, 2015b).

The EU Classification and Labelling Guidelines (ECHA 2012) uses a system based on the strength of evidence for hazard-identification as a "carcinogen". The guidelines allow for a chemical to be classified as not a carcinogen if a mode of action can be established to be not relevant to humans.

However, for carcinogens that are considered relevant for humans, the system does not distinguish between those which have a mode of action indicating that there is a presumption of no threshold for the dose response and those which have a mode of action indicating that there is a threshold for the dose response.

Categorization of carcinogenicity of agents (e.g., commodity or pesticide chemicals, food additives, viruses, or natural products) by the strength of evidence (e.g., animal cancer bioassays, epidemiology, other experimental in vitro and in vivo) without consideration of mode of action, dose-response and human exposure can result in agents being placed into the same category that vary widely in their likelihood to cause cancer. It has been suggested that the EU GHS process could be improved by including potency in a weight of evidence approach using methods which are already part of the EU Classification, Labeling and Packaging guidelines relating to the presence of substances classified as carcinogens in mixtures and preparations (Hennes et al, 2014).

10. Problems Resulting From Use of Outmoded Processes

The problems caused by hazard-identification classification schemes are complex and they have consequences for many parts of society. The original intent of these schemes was to identify chemicals or other agents which may be of concern and thus require further evaluation including full risk assessments to determine if action would be needed to mitigate a risk. We see this process working in the strategy used by the EPA (EPA, 2005) and UK Committee on Carcinogenicity (CoC, 2012) where the initial hazard-identification as a carcinogen triggers a logical and scientific consideration of the risk to human health which takes into account the potency, the mode of action and the magnitude, duration, frequency and route of exposure. Appropriate risk mismanagement decisions can then be made which can range from taking no action, using personal protective

equipment, decreasing personal exposure, restrictions on use, to outright banning of use in extreme circumstances.

However, all too often the public response to the classification by hazard-identification alone is not so reasoned. For example IARC classification as “carcinogenic to humans”, “probably carcinogenic to humans” or “possibly carcinogenic to humans” can all lead to negative publicity and a “health scare”. The “health scare” can trigger anxiety and lead to behavior which is detrimental to actually achieving desirable public health goals (Berry, 2016). Government and other agencies then have to use precious resources to respond because of publicly perceived rather than actual threats.

Several organisations have had to explain to the public what IARC does in attempts to alleviate unnecessary concern (Health Canada, 2016, Cancer Research UK, 2012). As an example, Cancer Research UK sums up what IARC does: *“Just because something is in IARC’s top level category, it doesn’t necessarily mean it’s public health number one – it’s more complex than that. IARC does ‘hazard identification’, not ‘risk assessment’. That sounds quite technical, but what it means is that IARC isn’t in the business of telling us how potent something is in causing cancer – only whether it does so or not. To take an analogy, think of banana skins. They definitely can cause accidents – but in practice this doesn’t happen very often (unless you work in a banana factory). And the sort of harm you can come to from slipping on a banana skin isn’t generally as severe as, say, being in a car accident. But under a hazard identification system like IARC’s, ‘banana skins’ and ‘cars’ would come under the same category – they both definitely do cause accidents.”*

Cancer UK (2012) explains that *“IARC categories are designed to flag things up to policy makers, so they can then analyse the scale of the problem, weigh the risks against the benefits, and bring in appropriate legislation.”* IARC sometimes assesses chemicals after they have already been considered in detail for both potential hazard and risk by stringent agencies responsible for

regulation, such as US EPA, EFSA, ECHA, JMPR, PMRA. This lack of coordination and co-operation can lead to problems, confusion, duplication of efforts, and the expenditure of unnecessary resources. A recent example of this has arisen with the herbicide glyphosate. The European Food Safety Authority has completed an extensive review of its original evaluation following a second mandate from the European Commission (EFSA, 2015) to consider the findings from IARC's classification of glyphosate as "*a probable human carcinogen*" (IARC, 2015). They came to the conclusion that "*glyphosate is unlikely to pose a carcinogenic hazard to humans and the evidence does not support classification with regard to its carcinogenic potential*" (EFSA 2015). Oddly, IARC reviewers have chosen to dispute the findings of the EFSA process and criticize EFSA for "*an over-reliance on non-publicly available industry-provided studies using a limited set of assays that define the minimum data necessary for the marketing of a pesticide*" (Portier et al 2016). In the case of pesticides, it is the legal responsibility of the regulated industry to provide all the studies needed to establish the safety of their products. Extensive data are legally mandated by regulatory authorities to enable them to evaluate efficacy and safety, and these data are generated in accordance with Good Laboratory Practice and a series of internationally harmonized and scientifically peer-reviewed study protocols, designed to maintain a high standard of scientific quality and consistency, and to provide confidence that study results are repeatable and acceptable. So it should not be surprising that there are large databases of studies sponsored by pesticides registrants given the legal and regulatory requirements by authorities, who have access to all the raw data. It is noteworthy that based on an weight of evidence approach that evaluates the consistency, dose response, time course, and biological plausibility of all relevant evidence, JMPR concluded that glyphosate "*is unlikely to pose a carcinogenic risk to humans from exposure through the diet*" (JMPR, 2016). EPA recently articulated its conclusions that "*The strongest support is for "not likely to be carcinogenic to humans at doses relevant to human health risk assessment."*" (EPA 2016)

The EU classification system can also cause problems because there are automatic risk management consequences built into downstream regulations. These are exemplified in the EU Directive on the regulation of crop protection products (EU, 2009). This regulation applies so-called cut off criteria which do not allow any products categorized as category 1A or 1B carcinogens to be registered for use. This is an example of a process where hazard-identification goes directly to risk management without going through hazard characterization and risk assessment, even though the regulations demand that a full toxicological and exposure data set be produced and a risk assessment be performed for every requested use. In many cases, a risk assessment using the modern approaches would show that different modes of action have different implications for human safety and would therefore impact the regulatory decision.

Chemicals and other agents, particularly those that are in widespread use that are flagged by classification processes based on hazard-identification alone tend to come under close scrutiny and become the subject of debate and concerted public campaigns. For instance the IARC classification of glyphosate has given rise to headlines such as “War in Europe- Battle over Glyphosate” (Genetic Literacy Project, 2016), “EU scientists in row over safety of Glyphosate weedkiller” Guardian (2016), and “How the World Health Organization’s cancer agency confuses consumers” (Reuters, 2016). The listing of a chemical by these hazard-identification only based schemes can have major implications either because of downstream regulation or because of the reputational damage which can be caused as a consequence of a distorted and misled public perception. Useful chemicals can be lost by attrition even without regulatory intervention, because of unfounded changes in public behavior and precautionary reaction of industry. In many cases, the categorization of a chemical by hazard-identification alone diverts effort and funding into research projects which continue long after the chemical has been determined not to pose a risk to humans or effective risk management actions have been taken. For instance, the NCI bioassay on chloroform was published in 1976 (NCI, 1976)

but over 90 papers have been published concerning its carcinogenicity (PubMed 2016). As recently as 2010, Take et al (2010) investigated interactions in kinetics of chloroform via the oral and inhalation routes in an attempt to put the results of the original high dose oral dosing based bioassay into the context of human exposure which is mainly by inhalation or via the dermal route (Take et al, 2010).

11. Problem Formulation: What Problem is Being Addressed?

“The mere formulation of a problem is far more essential than its solution, which may be merely a matter of mathematical or experimental skill. To raise new questions, new possibilities, to regard old problems from a new angle require creative imagination and marks real advances in science.” – Albert Einstein (1938).

Formulating the problem being addressed is key to solving it. Which problems are the hazard-identification based classification systems trying to address?

In the 1970s the problem could have been formulated as “identify those chemicals which are capable of causing cancer so they can be eliminated from use”. The processes put into place based on this problem formulation were set up in favor of identifying carcinogens, i.e. set up to minimize false negatives. The problem was formulated with the assumption that there would be a relatively small number of carcinogens which could be detected with reasonable reliability and did not foresee that there would turn out to be various mechanisms of carcinogenicity with different implications for risk assessment and risk management, and that a number of these would be specific to rodents. Most of the chemicals identified up to that time were potent DNA-reactive (genotoxic) carcinogens that produced cancer in both rodent models and in humans, such as aromatic amines, polycyclic aromatic hydrocarbons, nitrosamines and aflatoxins. These were believed to not have a threshold. With the development of the two year rodent bioassay incorporating an MTD, numerous chemicals

were identified as carcinogenic that were non-DNA reactive (non-genotoxic). Their cancer modes of action considered to be secondary consequence of their toxicity (e.g. sustained cytotoxicity or cell proliferation) and were considered to have a threshold. Subsequent research has shown that many produce cancer in rodents by a mode of action not relevant to humans. If relevant to humans, the presence of a threshold presents a completely different dose response and risk assessment than the DNA reactive carcinogens. Hazard-based systems do not distinguish these, even though potency and human risk are vastly different.

Hazard-identification based classification systems could have a part to play in addressing an updated problem formulation as an early warning or priority setting measure. There could be value if the processes used were rapid and were undertaken before more detailed risk assessment is completed and if they were acknowledged to provide a preliminary evaluation which would subsequently be refined as necessary. However, the current hazard-identification based systems are not treated as preliminary assessments leading to the problems which have been described in section 10.

Systems which are designed to place as many chemicals as possible into the most severe category are ultimately self-defeating. At first sight this could appear to be a benefit. More chemicals will be classified in the most severe category and would be subjected to stringent risk management, which in a hazard-identification only system means no exposure. This could be achieved at a fraction of the cost simply by assuming that all chemicals are carcinogens. But placing more and more chemicals into the most extreme category will have a severe unintended effect. In fact, when only the most hazardous chemicals are identified, the classification is respected and appropriate risk management decisions are taken, especially in those sectors where banning or withdrawal is not mandatory. However, if too many chemicals are placed into the most hazardous category, including those which do not represent an extreme risk, the distinction is lost, and respect for the system is eroded (American Cancer Society, 2016).

Many sectors will find it hard to operate by excluding all chemicals in this category and chemicals truly hazardous by adverse effects other than carcinogenicity may not be excluded or managed appropriately, thereby having the opposite effect from the intended one. For example, many chemicals that are natural components of various foods produce cancer in rodents at high doses, including substances in fruits and vegetables, which most people consider positive enhancers to health (Ames and Gold, 1990). Likewise, more than half of currently approved prescription pharmaceuticals are carcinogenic in rodent bioassay, and yet, based on a risk assessment rather than a hazard-based analysis, they benefit millions of grateful patients, with little or no risk of cancer (Brambilla et al, 2012).

It is now time to revise the problem formulation statement along the lines of “identify and characterize the carcinogenic potential of chemicals so that appropriate risk management measures can be taken to safeguard human health.”

This is the direction being taken by several national and international initiatives (e.g., Health Canada 2000; WHO IPCS 2010; EFSA 2013). The US EPA Framework for Human Health Risk Assessment to Inform Decision Making (EPA, 2014) outlines a stepwise approach that goes through Planning and scoping; Problem formulation; Risk assessment; Exposure and effects assessment (including hazard-identification and dose response assessment); Risk characterization; Public, stakeholder and community involvement; Informing decisions. The RISK21 process (Embry et al, 2014) emphasizes the importance of problem formulation, the use of existing data, and tiered assessment of exposure and then hazard in safety assessment.

Much has been achieved by international co-operation in advancing risk assessment it is now time for the categorization of carcinogenicity to be the subject of such an initiative. The WHO International Programme for Chemical Safety last reviewed classification schemes in 1995 (IPCS, 1995) and recognized the need to consider a range of issues which would need resolution to solve the problem of diverging classification schemes. These issues have been clarified in the intervening 20 years, and it is now the time to heed the call in the Lancet (Lancet, 2016) for an international initiative to develop a standardized, internationally agreed upon methodology for carcinogen assessment, coupled with tools for presenting results that are easily understood and accepted by all interested parties.

12. Reproductive Toxicology and Endocrine Disruption

This paper has focused on cancer but there are other areas of toxicology in which hazard-identification based classification systems exist, with similar untoward consequences. The classification system for reproductive and developmental toxicity in the EU is also a hazard-identification based system (Hennes et al, 2014). Classification as a Category 1A (known human reprotoxicant) or 1B (presumed human reprotoxicant) will trigger downstream automatic extreme risk management measures for some uses of chemicals, for instance plant protection products (EU 2009) using the so-called “cut-off” criteria. This causes similar problems to those caused by the hazard-identification based system for carcinogenicity within the EU. For example, vitamin A at high doses is a known human teratogen (Rothman et al, 1995). In a hazard-based classification system it should be banned, and yet lower exposures are essential for life. These problems are being reviewed in a Consultation on the Regulatory Fitness of Chemicals Legislation (EU, 2016).

Recently the EU published the proposed criteria for identifying chemicals which have the potential to cause adverse effects via endocrine disruption, but the proposed classification does not include an assessment of potency (EU 2016, a). It does not seem to be sensible to introduce a new hazard-

identification based system with all the problems that entails at the same time as the EU is reviewing the impact of such systems (EU 2016b).

13. Conclusions

Hazard-identification based classification schemes are inadequate to guide appropriate risk management decisions and have become outmoded. They are based on a concept developed in the 1970s that chemicals could be categorically divided cleanly into two classes; carcinogens and non-carcinogens. It was postulated at that time that a major reduction in cancer incidence would result if carcinogens could be identified and then eliminated from use and from the environment (Rettig, 2005). The classification that these schemes provide is based on the strength of evidence that the chemical has some degree of carcinogenic potential in humans or rodents but they do not indicate the degree of risk following real human exposure.

Categorization by the strength of evidence that a chemical has caused cancer in humans or in laboratory animals can place chemicals and agents with widely differing potencies in their ability to cause cancer and with very different modes of action into the same category. Chemicals with seven orders of magnitude difference in the dose required to cause cancer can be placed in the same category. This is how eating processed meat can fall into the same category as sulfur mustard gas.

More modern strategies based on problem formulation and hypothesis-based approaches in a risk decision framework such as those described by the US EPA (EPA, 2005, EPA, 2014), the UK Committee on Carcinogenicity (CoC, 2012), the EU Scientific Committee on Occupational Exposure Limits (Bolt and Huici-Montagud, 2008) provide clearer guidance and allow informed risk

management decisions to be taken. Once carcinogenic potential has been identified, hazard characterization then examines other factors such as the dose response and the mode of action to be combined with exposure assessment leading to risk assessment. Only then can risk management actions be taken if appropriate.

The hazard and risk characterization approach avoids the unintended downsides of creating health scares, incurring unnecessary economic costs and the diversion of public funds, which could be spent more wisely, into unnecessary research.

An international initiative to agree upon a standardized, internationally acceptable methodology for carcinogen assessment is needed now. The approach taken should incorporate principles and concepts of existing international consensus-based frameworks including the WHO IPCS mode of action framework.

Declaration of Interests

This work did not receive any specific support from funding agencies in the public, commercial, or not-for-profit sectors.

The authors' affiliations are as shown on the cover page. The authors had sole responsibility for the writing and content of the paper. The views and opinions expressed in the paper are those of the authors, and do not necessarily reflect the views or policies of the authors' current or former employers.

The authors have served as members of the following panels or committees and/or for the following organisations:

Council of Canadian Academies (VD); European Food Safety Authority (AB, AM); European Centre for Ecotoxicology and Toxicology of Chemicals (AB, JD); EU Scientific Committee on Occupational Exposure Limit Values (AM); European Medicines Agency (AB); Health Canada (PF-C); International Agency for Research on Cancer (SC, DW); International Life Science Institute (AB, SC, JD, VD, PF-C, AM, JS, RS, DW); Joint WHO/FAO Meeting on Pesticides Residues (AB, VD, PF-C); Italian Committee on Pesticides (AM); Joint WHO/FAO Expert Committee on Food Additives (Residues of Veterinary Drugs) (AB); National Institutes of Health (SC); National Academy of Sciences (SC, PF-C, RS); National Institute of Environmental Health Sciences (SC, DW); National Toxicology Program (SC, VD, PF-C, JS, DW); Organization for Economic Cooperation and Development (VD, PF-C, JS, RS); Swiss Centre for Applied Human Toxicology (AB, AM); UK Advisory Committee on Pesticides (AB); United Kingdom Committee on Carcinogenicity (AB, JD); UK Committee on the Medical Effects of Air Pollutants (AB); UK Committee on Residues of Veterinary Drugs (AB); UK Committee on Toxicity (AB); United States Environmental Protection Agency (SC, VD, PF-C, JS, RS, DW); United States Food and Drug Administration (SC, PF-C); World Health Organization International Program on Chemical Safety (AB, SC, JD, VD, PF-C, AM, JS, RS).

VD is retired from the US Environmental Protection Agency.

PF-C is retired from the US Environmental Protection Agency and the International Life Sciences Institute.

JS is retired from the US Environmental Protection Agency.

RS is retired from the US Environmental Protection Agency.

References

ACGIH (2016) Documentation of the Threshold Limit Values and Biological Exposure Indices, 7th Ed Supplement

American Cancer Society (2016)

<http://www.cancer.org/cancer/cancercauses/othercarcinogens/athome/cancer-warning-labels-based-on-californias-proposition-65>; accessed Oct 9, 2016

Ames BN and Gold LW (1990) Chemical carcinogenesis: Too many rodent carcinogens? *Proc Natl Acad Sci* 87: 7272-7276

Barlow SM, Boobis AR, Bridges J, Cockburn A, Dekant W, Hepburn P, Houben GF, Konig J, Nauta MJ, Schuermans J, Banati D (2015) The role of hazard- and risk-based approaches in ensuring food safety. *Trends in Food Science & Technology* 46, 176-188

Berry C (2016) The Dangers of Hazards *Toxicol. Res.*, 2016, 5, 373

Bolt HM and Huici-Montagud A (2008) Strategy of the scientific committee on occupational exposure limits (SCOEL) in the derivation of occupational exposure limits for carcinogens and mutagens. *Archives Toxicology* 82(1):61-64

Boobis A, Cohen S, Dellarco V, McGregor D, Meek M, Vickers C, Willcocks D, Farland W (2006) IPCS framework for analyzing the relevance of a cancer mode of action for humans. *Crit Rev Toxicol.* **36**, 781-92.

- Boobis, AR, Doe, JE, Heinrich-Hirsch, B, Meek, ME, Munn, S, Ruchirawat, M, Schlatter, J, Seed, J, Vickers, C. (2008). IPCS framework for analyzing the relevance of a noncancer mode of action for humans. Crit Rev Toxicol 38, 87-96.
- Brambilla, G, Mattioli, F, Robbiano, L, Martelli, A. (2012). Update of carcinogenicity studies in animals and humans of 535 marketed pharmaceuticals. Mutat Res 750(1): 1-51.
- Cancer Research UK (2012) <http://scienceblog.cancerresearchuk.org/2012/06/14/diesel-fumes-definitely-cause-cancer-should-we-be-worried/> Accessed 13 October 2016
- CoC (2012) A guidance statement from the Committee on Carcinogenicity of Chemicals in Food, Consumer Products and the Environment (COC) - A Strategy for the Risk Assessment of Chemical Carcinogens COC/G1 – Version 4 (2012)
- ECHA (2012). Guidance on the application of the CLP criteria. Guidance to Regulation (EC) No 1272/2008 on classification, labelling and packaging (CLP) of substances and mixtures. Version 3.0 November 2012.
- EFSA. 2013. International frameworks dealing with human risk assessment of combined exposure to multiple chemicals. EFSA J 11(7):3313, doi:10.2903/j.efsa.2013.3313.
- EFSA (2015) Conclusion on the peer review of the pesticide risk assessment of the active substance glyphosate EFSA Journal 2015;13(11):4302

747

748 Einstein (1938) in The Evolution of Physics, by A Einstein and L Infeld, published by Simon and
749 Schuster, New York, p95

750

751 Embry MR, Bachman AN, Bell DR, Boobis AR, Cohen SM, Dellarco M, Dewhurst IC, Doerr NG, Hines
752 RN, Moretto A, Pastoor TP, Phillips RD, Rowlands JC, Tanir JY, Wolf DC, Doe JE (2014) Risk
753 Assessment in the 21st Century: Roadmap and Matrix Critical Reviews in Toxicology. 2014;44(Suppl
754 3):6-16

755

756 EPA (2005) Guidelines for Carcinogen Risk Assessment EPA/630/P-03/001F

757

758 EPA (2014) Framework for Human Health Risk Assessment to Inform Decision Making April 5, 2014

759 U.S. Environmental Protection Agency Office of the Science Advisor Risk Assessment Forum

760

761 EPA (2015) Chemicals Evaluated for Carcinogenic Potential; Office of Pesticide Programs, U.S.

762 Environmental Protection Agency; Annual Cancer Report 2015

763

764 EPA 92016) Glyphosate Issue Paper: Evaluation of Carcinogenic Potential EPA's Office of Pesticide
765 Programs September 12, 2016

766

767 JOINT FAO/WHO MEETING ON PESTICIDE RESIDUES Geneva, 9–13 May 2016 SUMMARY REPORT

768

769 EU (2009) Regulation (EC) No 1107/2009 of the European Parliament and of the Council of
770 21 October 2009 concerning the placing of plant protection products on the market and repealing
771 Council Directives 79/117/EEC and 91/414/EEC
772
773 EU (2014) Defining criteria for identifying Endocrine Disruptors in the context of the implementation
774 of the Plant Protection Product Regulation and Biocidal Products Regulation
775 [http://ec.europa.eu/smart-](http://ec.europa.eu/smart-regulation/impact/planned_ia/docs/2014_env_009_endocrine_disruptors_en.pdf)
776 [regulation/impact/planned_ia/docs/2014_env_009_endocrine_disruptors_en.pdf](http://ec.europa.eu/smart-regulation/impact/planned_ia/docs/2014_env_009_endocrine_disruptors_en.pdf) Accessed 13 Oct
777 2016
778
779 EU(2016a) setting out scientific criteria for the determination of endocrine disrupting properties and
780 amending Annex II to Regulation (EC) 1107/2009
781 http://ec.europa.eu/health/endocrine_disruptors/docs/2016_pppcriteria_en.pdf Accessed 13 Oct
782 2016
783
784 EU (2016b) DG Growth Consultation on the regulatory fitness of chemicals legislation (excluding
785 REACH) http://ec.europa.eu/growth/tools-databases/newsroom/cf/itemdetail.cfm?item_id=8695
786 Accessed 13 Oct 2016
787
788 GBD2010 (2012) A comparative risk assessment of burden of disease and injury attributable to 67
789 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global
790 Burden of Disease Study 2010, Lancet 380 2224-2260
791

792 Genetic Literacy Project, 2016 <https://www.geneticliteracyproject.org/2016/08/16/war-europe->
793 [battle-glyphosate](#) Accessed 13 Oct 2016

794

795

796 Gold L, Slone T and Bernstein L (1989) Summary of Carcinogenic Potency and Positivity for 492
797 Rodent Carcinogens in the Carcinogenic Potency Database. Environmental Health Perspectives, 79,
798 259-272

799

800 Goldblatt M (1944) Investigation of Toxic Hazards. Br J Ind Med 1944; 1 20-30

801

802 Guardian (2016) EU scientists in row over safety of Glyphosate weedkiller 13 January 2016
803 <https://www.theguardian.com/environment/2016/jan/13/eu-scientists-in-row-over-safety-of->
804 [glyphosate-weedkiller](#) Accessed 13 Oct 2016

805

806 Health Canada. 2000. Health Canada Decision-Making Framework for Identifying, Assessing,
807 and Managing Health Risks. Ottawa, Ontario, Canada:Health Canada.

808 http://www.hc-sc.gc.ca/ahc-asc/alt_formats/hpfb-dgpsa/pdf/pubs/risk-risques-eng.pdf
809 Accessed 13 Oct 2016

810

811 Health Canada (2016) http://www.hc-sc.gc.ca/cps-spc/pest/part/consultations/_prvd2015-
812 [01/prvd2015-01-eng.php](#) Accessed 13 Oct 2016

813

814 Hennes C, Batke M, Bomann W, Duhayon S, Kosemund K, Politano V, Stinchcombe S, Doe J (2014)

815 Incorporating potency into the EU classification for carcinogenicity and reproductive

816 toxicity Regulatory Toxicology and Pharmacology, 70 457–67
817
818 Hill, AB (1965). The Environment and Disease: Association or Causation. Proceedings of the Royal
819 Society of Medicine 58 295-300
820
821 Hueper W (1955) Lung Cancers and their Causes CA: A Cancer Journal for Clinicians 5 95–100
822
823 IARC (2012) IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Some
824 Organophosphate Insecticides and Herbicides: Diazinon, Glyphosate, Malathion, Parathion, and
825 Tetrachlorvinphos Volume 112
826
827 IARC (2015a) IARC Monographs on the Evaluation of Carcinogenic Risk to Humans: Preamble. WHO
828 IARC updated 2015
829
830 IARC (2015b) Q&A on the carcinogenicity of the consumption of red meat and processed meat
831 https://www.iarc.fr/en/media-centre/iarcnews/pdf/Monographs-Q&A_Vol114.pdf Accessed 13 Oct
832 2016
833
834 IPCS (1995) IPCS Workshop on the Harmonization of Risk Assessment for Carcinogenicity and
835 Mutagenicity (Germ Cells) – A Scoping Meeting Carshalton UK, February 1995
836
837 JMPR (2016) JOINT FAO/WHO MEETING ON PESTICIDE RESIDUES Geneva, 9–13 May 2016 SUMMARY
838 REPORT
839

- 840 Julien E, Boobis A, Olin S and ILSI Research Foundation Threshold Working group (2009) The Key
841 Events Dose-Response Framework: A Cross-Disciplinary Mode-of-Action Based Approach to
842 Examining Dose-Response and Thresholds *Crit Rev Food Sci Nutr* 49, 682-689.
- 843
- 844 Kauppinen T, Uuksulainen S, Saalo A, Makinen I and Pukkala E (2013) Use of the Finnish Information
845 System on Occupational Exposure (FINJEM) in Epidemiologic, Surveillance, and Other Applications
846 *Ann Occup Hyg* 57:593-609
- 847
- 848 Krewski et al (2010) Toxicity Testing in the 21st Century: A Vision and a Strategy *J Toxicol and*
849 *Environ Health B Crit Rev* 13 51-138
- 850
- 851 *Lancet* (2016) When is a Carcinogen not a Carcinogen? *Lancet Oncology* 17, 681
- 852
- 853 Meek, ME, Bucher, JR, Cohen, SM, Dellarco, V, Hill, RN, Lehman-McKeeman, LD, Longfellow, DG,
854 Pastoor, T, Seed, J, Patton, DE.. (2003). A framework for human relevance analysis of information on
855 carcinogenic modes of action. *Crit Rev Toxicol* **33**, 591-653.
- 856
- 857 Meek ME, Boobis A, Cote I, Dellarco V, Fotakis G, Munn S, Seed J, Vickers C. (2014) New
858 developments in the evolution and application of the WHO/IPCS framework on mode of
859 action/species concordance analysis *J Appl Toxicol* 34 1-18.
- 860
- 861 NCI (1976) Report on the carcinogenesis Bioassay of Chloroform (CAS No 67-66-3) National
862 Toxicology Program National Cancer Institute Technical Report Series March 1976 1-60

863

864 OECD (2009) OECD Guidelines for the testing of Chemicals, Section 4, Health Effects, Test No 451:
865 Carcinogenicity Studies.

866

867 Pastoor T and Stevens J (2005) Historical perspective of the cancer bioassay. Scand. J. Work Environ.
868 Health 2005;31 Suppl. 1:129-140.

869

870 Portier, C et al (2015) Differences in the carcinogenic evaluation of glyphosate between the
871 International Agency for Research on Cancer (IARC) and the European Food Safety Authority (EFSA)
872 J Epidemiol Community Health doi:10.1136/jech-2015-207005

873

874 PubMed (2016) <http://www.ncbi.nlm.nih.gov/pubmed> search using term 'chloroform
875 carcinogenicity' Accessed 13 Oct 2016

876

877 Rettig, R (2007) Cancer Crusade – The Story of the National Cancer Act of 1971 Authors Choice Press
878 ISBN-13:978-0-595-35847-2

879

880 Reuters (2016) How the World Health Organization's cancer agency confuses consumers.
881 <http://www.reuters.com/investigates/special-report/health-who-iarc/> Accessed 13 Oct 2016

882

- 883 Roser M (2015) – ‘Life Expectancy’. Published online at OurWorldInData.org. Retrieved from:
884 <http://ourworldindata.org/data/population-growth-vital-statistics/life-expectancy/> Accessed 13 Oct
885 2016
886
- 887 Rothman KJ, Moore LL, Singer MR, Nouven US, Mannino S, Milunsky A(1995) Teratogenicity of high
888 vitamin A intake. N Engl J Med 333, 1369-73.
889
- 890 Seed, J, Carney, EW, Corley, RA, Crofton KM, DeSesso, JM, Foster, PM, Kavlock, R, Kimmel, G,
891 Klaunig, J, Meek, ME, Preston, RJ, Slikker, W, Jr., Tabacova, S, William, GM (2005). Overview: Using
892 Mode of Action and Life Stage Information to Evaluate the Human Relevance of Animal Toxicity
893 Data. Crit Rev Toxicol 35, 663-672.
894
- 895 Simon TW, Stoney Simons S, Preston RJ Boobis A, Cohen SM, Doerrner NG, Fenner-Crisp PA, McMullin
896 CA, McQueen CA, Rowlands JC (2014)The use of mode of action information in risk assessment:
897 Quantitative key events/dose-response framework for modeling the dose-response for key events
898 Critical Reviews In Toxicology 44 , ,2014 Supp 3 17-43
899
- 900 Slikker W Jr, Andersen ME, Bogdanffy MS, Bus JS, Cohen SD, Conolly RB, David RM, Doerrner NG,
901 Dorman DC, Gaylor DW, Hattis D, Rogers JM, Setzer RW, Swenberg JA, Wallace K. 2004a. Dose-
902 dependent transitions in mechanisms of toxicity. Toxicol Appl Pharmacol. 2004 Dec 15;201(3):203-
903 25. Review.
904
- 905 Slikker W Jr, Andersen ME, Bogdanffy MS, Bus JS, Cohen SD, Conolly RB, David RM, Doerrner NG,
906 Dorman DC, Gaylor DW, Hattis D, Rogers JM, Setzer RW, Swenberg JA, Wallace K. 2004b. Dose-

907 dependent transitions in mechanisms of toxicity: case studies. Toxicol Appl Pharmacol. 2004 Dec
 908 15;201(3):226-94. Review.

909

910 Sonich-Mullin C, Fielder R, Wiltse J, Baetcke K, Dempsey J, Fenner-Crisp J, Grant D, Hartley M,
 911 Knaap A, Kroese D, Mangelsdorf I, Meek E, Rice J, and Younes M (2001) IPCS Conceptual
 912 Framework for Evaluating a Mode of Action for Chemical Carcinogenesis Regulatory Toxicology and
 913 Pharmacology 34, 146–152

914

915 Take M, Yamamoto S, Ohnishi M, Matsumoto M, Nagano K, Nagato K, Hirota T, Fukushima S (2010)
 916 Chloroform distribution and accumulation by combined inhalation plus oral exposure routes in rats. J
 917 Environ Sci Health A Tox Hazard Subst Environ Eng 45, 1616-1624

918

919 UN (2011) Globally Harmonised System of Classification and Labelling of Chemicals (GHS) Fourth
 920 Revised Edition, United Nations, Foreword Section 2.

921

922 WHO IPCS. World Health Organisation International Programme on Chemical Safety. 2010. WHO
 923 Human Health Risk Assessment Tool Kit: Chemical Hazards. Geneva:WHO. Available:
 924 <http://www.who.int/ipcs/publications/methods/harmonization/toolkit.pdf?ua=1> Accessed 13
 925 October 2016